

UC San Diego

UC San Diego Previously Published Works

Title

Draft Genome Sequence of Rhodococcus sp. Strain ATCC 49988, a Quinoline-Degrading Bacterium.

Permalink

<https://escholarship.org/uc/item/11x2j0s7>

Journal

Microbiology resource announcements, 8(25)

ISSN

2576-098X

Authors

Gupta, Nidhi
Skinner, Kelly A
Summers, Zarath M
et al.

Publication Date

2019-06-01

DOI

10.1128/mra.00403-19

Peer reviewed



Draft Genome Sequence of *Rhodococcus* sp. Strain ATCC 49988, a Quinoline-Degrading Bacterium

Nidhi Gupta,^{a,b} Kelly A. Skinner,^a Zarath M. Summers,^c Janaka N. Edirisinghe,^{a,b} José P. Faria,^{a,b} Christopher W. Marshall,^{a*} Anukriti Sharma,^{b*} Neil R. Gottel,^{b*} Jack A. Gilbert,^{a,b*} Christopher S. Henry,^{a,b} Edward J. O'Loughlin^a

^aArgonne National Lab (ANL), Argonne, Illinois, USA

^bUniversity of Chicago, Chicago, Illinois, USA

^cExxonMobil Research and Engineering Company, Annandale, New Jersey, USA

ABSTRACT We report here the 4.9-Mb genome sequence of a quinoline-degrading bacterium, *Rhodococcus* sp. strain ATCC 49988. The draft genome data will enable the identification of genes and future genetic modification to enhance traits relevant to heteroaromatic compound degradation.

Quinoline is a nitrogen-containing heterocyclic compound that is used as a solvent in the chemical industry and occurs widely in coal tar, oil shale, and plant alkaloids. It is known to be a mutagenic and carcinogenic compound (1). Due to its widespread use and toxic properties, the degradation of quinoline by microorganisms has been studied extensively. *Rhodococcus* sp. strain ATCC 49988 was isolated from soil by enrichment culture using quinoline as a dominant carbon, nitrogen, and energy source (2). In order to understand the genetic basis of quinoline degradation and other abilities of this strain, genome sequence analysis of *Rhodococcus* sp. strain ATCC 49988 was carried out.

Rhodococcus sp. strain ATCC 49988 was obtained from the ATCC. It was grown in tryptic soy broth at 30°C with shaking at 200 rpm for 16 h, after which DNA for whole-genome sequencing was isolated using a DNeasy PowerSoil kit (catalog number 12888-50; Qiagen), followed by preparation of an Illumina Nextera XT paired-end library. Whole-genome sequencing was performed using a MiSeq system (Illumina) and produced 13,615,724 (average length, 160 bp) paired-end reads. The raw sequence was processed using the KBase (3) platform (<https://narrative.kbase.us/narrative/ws.39490.obj.1>), and the sequence was uploaded in FASTQ format. During analysis, default parameters were used for all software, unless otherwise specified. The read quality was assessed using FastQC v 0.11.5 (4), and the adaptor sequences specific to the Nextera DNA library were trimmed using Trimmomatic v 0.36 (5). *De novo* assembly conducted using SPAdes v 3.12.0 (6) yielded 27 contigs. The draft genome sequence has a total length of 4,970,306 bp (N_{50} , 792,857 bp) and a GC content of 67.87%. The genome coverage was 155.0×. Genome assembly statistics were computed using QUAST v 0.0.4 (7). The assembled contigs ranged in size from 805 bp to 1,052,617 bp. The assembly was evaluated for completeness and contamination using CheckM v 1.0.8 (8), which characterized the assembly as 99.4% complete with 0% contamination.

The genome sequence of this *Rhodococcus* sp. strain was first annotated using Prokka v 1.12 (9), which predicted a total of 4,508 protein-coding genes in the genome (<https://narrative.kbase.us/#dataview/39490/15/1>). Next, the genome was functionally annotated using Rapid Annotations using Subsystems Technology (RAST) v 0.1.1 (10; <https://narrative.kbase.us/#dataview/39490/17/1>), which assigned 1,406 (31%) of the genes to SEED subsystems. Several putative polycyclic aromatic hydrocarbon (PAH)

Citation Gupta N, Skinner KA, Summers ZM, Edirisinghe JN, Faria JP, Marshall CW, Sharma A, Gottel NR, Gilbert JA, Henry CS, O'Loughlin EJ. 2019. Draft genome sequence of *Rhodococcus* sp. strain ATCC 49988, a quinoline-degrading bacterium. Microbiol Resour Announc 8:e00403-19. <https://doi.org/10.1128/MRA.00403-19>.

Editor David Rasko, University of Maryland School of Medicine

Copyright © 2019 Gupta et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Edward J. O'Loughlin, oloughlin@anl.gov.

* Present address: Christopher W. Marshall, University of Pittsburgh, Pittsburgh, Pennsylvania, USA; Anukriti Sharma, Neil R. Gottel, and Jack A. Gilbert, University of California, San Diego, San Diego, California, USA.

Received 4 April 2019

Accepted 26 May 2019

Published 20 June 2019

catabolic enzymes (11), such as monooxygenase, dioxygenases, dehydrogenases, and cytochrome P450, were found in the *Rhodococcus* genome. However, genes for quinoline 2-oxidoreductase were not annotated in the genome, indicating that this species uses a different enzyme to convert quinoline to 2-hydroxyquinoline. The genomic information of this *Rhodococcus* sp. strain will not only facilitate our understanding of the metabolism for recalcitrant heteroaromatic compounds but will also expand our knowledge of the physiology of the *Rhodococcus* genus.

Data availability. The draft genome sequence of this *Rhodococcus* sp. strain has been deposited in GenBank under the accession numbers [SDMJ01000001](#) to [SDMJ01000026](#), and the raw sequencing reads are available in the Sequence Read Archive under accession number [SRP179964](#).

ACKNOWLEDGMENT

This work was supported by ExxonMobil via cooperative research and development agreement (CRADA) number C1400201.

REFERENCES

1. Felczak A, Zawadzka K, Lisowska K. 2014. Efficient biodegradation of quinolone—factors determining the process. *Int Biodeterior Biodegradation* 96:127–134. <https://doi.org/10.1016/j.ibiod.2014.08.004>.
2. O'Loughlin EJ, Kehrmeyer SR, Sims GK. 1996. Isolation, characterization and substrate utilization of a quinoline-degrading bacterium. *Int Biodeterior Biodegradation* 38:107–118. [https://doi.org/10.1016/S0964-8305\(96\)00032-7](https://doi.org/10.1016/S0964-8305(96)00032-7).
3. Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D, Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Chandonia J-M, Chia J-M, Colasanti R, Conrad N, Davis JJ, Davison BH, DeJongh M, Devoid S, Dietrich E, Dubchak I, Edirisinghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W, Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, et al. 2018. KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 36:566–569. <https://doi.org/10.1038/nbt.4163>.
4. Brown J, Pirrung M, McCue LA. 2017. FQC Dashboard: integrates FastQC results into a Web-based, interactive, and extensible FASTQ quality control tool. *Bioinformatics* 33:3137–3139. <https://doi.org/10.1093/bioinformatics/btx373>.
5. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
7. Mikheenko A, Valin G, Pribelski A, Saveliev V, Gurevich A. 2016. Icarus: visualizer for de novo assembly evaluation. *Bioinformatics* 32:3321–3323. <https://doi.org/10.1093/bioinformatics/btw379>.
8. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
9. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
10. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
11. Haritash AK, Kaushik CP. 2009. Biodegradation aspects of polycyclic aromatic hydrocarbons (PAHs): a review. *J Hazard Mater* 169:1–15. <https://doi.org/10.1016/j.jhazmat.2009.03.137>.